

MTH 1126 Test #3 (11am Class) - Solutions
SPRING 2022

Pat Rossi

Name _____

Show CLEARLY how you arrive at your answers.

1. $\int \cos^2(x) dx =$

We have $\sin(x)$ raised to an even power and no factor of $\cos(x)$. U-sub won't work here. We have to express $\sin^2(x)$ in a form other than something raised to a power.

We can do this by using the identity: $\cos^2(x) = \frac{1+\cos(2x)}{2}$

$$\text{Thus, } \int \cos^2(x) dx = \int \frac{1+\cos(2x)}{2} dx = \int \left(\frac{1}{2} + \frac{1}{2} \cos(2x)\right) dx = \int \frac{1}{2} dx + \frac{1}{2} \int \underbrace{\cos(2x)}_{\cos(u)} \underbrace{dx}_{\frac{1}{2} du}$$

$$= \frac{1}{2}x + \frac{1}{2} \left(\frac{1}{2} \sin(2x)\right) + C = \frac{1}{2}x + \frac{1}{4} \sin(2x) + C$$

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4} \sin(2x) + C$$

2. $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} =$

$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} \sim \frac{0}{0}$ (Use L'Hopital's Rule)

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}[\sin(x)]}{\frac{d}{dx}[x]} = \lim_{x \rightarrow 0} \frac{\cos(x)}{1} = \frac{1}{1} = 1$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

3. $\int \frac{3x+20}{x^2-3x-10} dx =$

Note that $\int \frac{3x+20}{x^2-3x-10} dx$ does not fit the form: $\int \frac{1}{u} du$

$$\begin{aligned} u &= x^2 - 3x - 10 \\ \frac{du}{dx} &= 2x - 3 \\ du &= (2x - 3) dx \end{aligned}$$

$$\int \frac{3x+20}{x^2-3x-10} dx = \int \underbrace{\frac{1}{x^2-3x-10}}_{\frac{1}{u}} \underbrace{(3x+20) dx}_{\text{NOT a constant multiple of } du}$$

Therefore, we decompose $\frac{3x+20}{x^2-3x-10}$ into the sum of simpler quotients:

1. Make sure that $\deg(\text{numerator}) \leq \deg(\text{denominator})$

2. Factor the denominator.

$$\frac{3x+20}{x^2-3x-10} = \frac{3x+20}{(x-5)(x+2)}$$

3. For each linear factor $(x + c)$, form the term $\frac{C_1}{x+c}$

$$\frac{3x+20}{x^2-3x-10} = \frac{3x+20}{(x-5)(x+2)} = \frac{C_1}{x-5} + \frac{C_2}{x+2}$$

4. Solve for the constants

$$\frac{3x+20}{(x-5)(x+2)} = \frac{C_1}{x-5} + \frac{C_2}{x+2}$$

$$\Rightarrow \frac{3x+20}{(x-5)(x+2)} (x-5)(x+2) = \frac{C_1}{x-5} (x-5)(x+2) + \frac{C_2}{x+2} (x-5)(x+2)$$

$$\text{i.e., } 3x + 20 = C_1(x + 2) + C_2(x - 5)$$

Plu in “strategic values” of x to find the values of the constants.

$$\boxed{x = -2}$$

$$\Rightarrow 14 = -7C_2$$

$$\Rightarrow \boxed{C_2 = -2}$$

$$\boxed{x = 5}$$

$$\Rightarrow 35 = 7C_1$$

$$\Rightarrow \boxed{C_1 = 5}$$

$$\text{Thus, } \frac{3x+20}{x^2-3x-10} = \frac{C_1}{x-5} + \frac{C_2}{x+2} = \frac{5}{x-5} - \frac{2}{x+2}$$

$$\text{i.e., } \frac{3x+20}{x^2-3x-10} = \frac{5}{x-5} - \frac{2}{x+2}$$

$$\text{Consequently, } \int \frac{3x+20}{x^2-3x-10} dx = \int \left(\frac{5}{x-5} - \frac{2}{x+2} \right) dx = 5 \int \frac{1}{x-5} dx - 2 \int \frac{1}{x+2} dx = 5 \ln |x - 5| - 2 \ln |x + 2| + C$$

$$\boxed{\int \frac{3x+20}{x^2-3x-10} dx = 5 \ln |x - 5| - 2 \ln |x + 2| + C}$$

4. $\int \sin^7(x) \cos^3(x) dx$ = We have odd powers of $\sin(x)$ and $\cos(x)$.

Therefore, we can use either $\sin(x)$ or $\cos(x)$ as our “future du.”

1. Reserve a factor of $\cos(x)$ to serve as our “future du.”

$$= \int \sin^7(x) \cos^2(x) \underbrace{\cos(x) dx}_{\text{“future du”}}$$

This means that we intend to let $u = \sin(x)$

2. Convert remaining cosines into sines

$$= \int \sin^7(x) \cos^2(x) \underbrace{\cos(x) dx}_{\text{“future du”}} = \int \sin^7(x) (1 - \sin^2(x)) \cos(x) dx = \int (\sin^7(x) - \sin^9(x)) \cos(x) dx$$

Let $u = \sin(x)$
$\Rightarrow \frac{du}{dx} = \cos(x)$
$\Rightarrow du = \cos(x) dx$

$$= \int \underbrace{(\sin^7(x) - \sin^9(x))}_{u^7 - u^9} \underbrace{\cos(x) dx}_{du} = \int (u^7 - u^9) (du) = \frac{u^8}{8} - \frac{u^{10}}{10} + C$$

$$= \frac{\sin^8(x)}{8} - \frac{\sin^{10}(x)}{10} + C$$

$\int \sin^7(x) \cos^3(x) dx = \frac{\sin^8(x)}{8} - \frac{\sin^{10}(x)}{10} + C$

Alternative Solution appears on the next page

$$\int \sin^7(x) \cos^3(x) dx =$$

We have odd powers of $\sin(x)$ and $\cos(x)$.

Therefore, we can use either $\sin(x)$ or $\cos(x)$ as our “future du.”

1. Reserve a factor of $\sin(x)$ to serve as our “future du.”

$$= \int \sin^6(x) \cos^2(x) \underbrace{\sin(x) dx}_{\text{“future du”}}$$

This means that we intend to let $u = \cos(x)$

2. Convert remaining sines into cosines

$$= \int (\sin^2(x))^3 \cos^2(x) \sin(x) dx = \int (1 - \cos^2(x))^3 \cos^2(x) \sin(x) dx$$

$$= \int (-\cos^6 x + 3\cos^4 x - 3\cos^2 x + 1) \cos^2(x) \sin(x) dx$$

$$= \int (-\cos^8 x + 3\cos^6 x - 3\cos^4 x + \cos^2(x)) \sin(x) dx$$

$$\begin{aligned} \text{Let } & u = \cos(x) \\ \Rightarrow & \frac{du}{dx} = -\sin(x) \\ \Rightarrow & du = -\sin(x) dx \\ \Rightarrow & -du = \sin(x) dx \end{aligned}$$

$$= \int (-u^8 + 3u^6 - 3u^4 + u^2) (-du) = \int (u^8 - 3u^6 + 3u^4 - u^2) (du)$$

$$= \frac{1}{9}u^9 - \frac{3}{7}u^7 + \frac{3}{5}u^5 - \frac{1}{3}u^3 + C$$

$$= \frac{1}{9} \cos^9(x) - \frac{3}{7} \cos^7(x) + \frac{3}{5} \cos^5(x) - \frac{1}{3} \cos^3(x) + C$$

$$\int \sin^7(x) \cos^3(x) dx = \frac{1}{9} \cos^9(x) - \frac{3}{7} \cos^7(x) + \frac{3}{5} \cos^5(x) - \frac{1}{3} \cos^3(x) + C$$

5. $\int x\sqrt{4-9x^2}dx =$ (Use Trig Substitution)

This can be done easily by U-substitution, but I think that we were instructed to do this using “Trig Substitution.”

We match the radical $\sqrt{4-9x^2}$ with the radical $\sqrt{a^2-a^2\sin^2(\theta)}$

\Rightarrow	$a^2 = 4$
\Rightarrow	$a = 2$
\Rightarrow	$9x^2 = a^2 \sin^2(\theta)$
\Rightarrow	$3x = a \sin(\theta)$
\Rightarrow	$x = \frac{a}{3} \sin(\theta)$
\Rightarrow	$\frac{dx}{d\theta} = \frac{a}{3} \cos(\theta)$
\Rightarrow	$dx = \frac{a}{3} \cos(\theta) d\theta$

Rewrite the integral in terms of θ


$$\int x\sqrt{4-9x^2}dx = \int \frac{a}{3} \sin(\theta) \sqrt{a^2-a^2\sin^2(\theta)} \frac{a}{3} \cos(\theta) d\theta = \int \frac{a}{3} \sin(\theta) \sqrt{a^2 \cos^2(\theta)} \frac{a}{3} \cos(\theta) d\theta$$

$$= \int \frac{a}{3} \sin(\theta) a \cos(\theta) \frac{a}{3} \cos(\theta) d\theta = \frac{a^3}{9} \int \underbrace{\cos^2(\theta)}_{u^2} \underbrace{\sin(\theta)}_{-du} d\theta$$

\Rightarrow	$u = \cos(\theta)$
\Rightarrow	$\frac{du}{d\theta} = -\sin(\theta)$
\Rightarrow	$du = -\sin(\theta) d\theta$
\Rightarrow	$-du = \sin(\theta) d\theta$

$$= \frac{a^3}{9} \int u^2 (-du) = -\frac{a^3}{9} \int u^2 du = -\frac{a^3}{9} \frac{u^3}{3} + C = -\frac{a^3 \cos^3(\theta)}{9} + C$$

To convert back to x , recall that $x = \frac{a}{3} \sin(\theta)$ (i.e., $\sin(\theta) = \frac{3x}{a} = \frac{\text{opp}}{\text{hyp}}$)



$$\int x\sqrt{4-9x^2} dx = \dots = -\frac{a^3 \cos^3(\theta)}{9 \cdot 3} + C = -\frac{a^3 \left(\frac{\sqrt{a^2-9x^2}}{a}\right)^3}{9 \cdot 3} + C = -\frac{a^3 \left(\frac{a^2-9x^2}{a^3}\right)^{\frac{3}{2}}}{9 \cdot 3} + C$$

$$= -\frac{1}{9} \frac{(a^2-9x^2)^{\frac{3}{2}}}{3} + C = -\frac{(4-9x^2)^{\frac{3}{2}}}{27} + C = -\frac{1}{27} (4-9x^2)^{\frac{3}{2}} + C$$

$$\int x\sqrt{4-9x^2} dx = -\frac{1}{27} (4-9x^2)^{\frac{3}{2}} + C$$

6. $\lim_{x \rightarrow 0} \frac{e^x - 1}{\ln(x+1)} \sim \frac{0}{0}$

Use L'Hopital's Rule!

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{\ln(x+1)} = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}[e^x - 1]}{\frac{d}{dx}[\ln(x+1)]} = \lim_{x \rightarrow 0} \frac{e^x}{\frac{1}{x+1}} = \frac{e^0}{\frac{1}{(0)+1}} = 1$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{\ln(x+1)} = 1$$

7. $\int e^x \sin(x) dx =$ (Use Integration by Parts)

When doing integration by parts and the integrand consists of two functions that have derivatives that are “cyclic,” the rule of thumb is that either function can be u and either function can be dv . We repeat the process of Integration by Parts until we get the original integral. Then we solve for the integral algebraically.

Arbitrarily, we'll let $u = e^x$ and $dv = \sin(x) dx$

This yields:

u	$=$	e^x	dv	$=$	$\sin(x) dx$
$\frac{du}{dx}$	$=$	e^x	$\int dv$	$=$	$\int \sin(x) dx$
du	$=$	$e^x dx$	v	$=$	$-\cos(x)$

Thus we have:

$$\begin{aligned} \int \underbrace{e^x}_u \underbrace{\sin(x) dx}_{dv} &= \int u dv = uv - \int v du = e^x (-\cos(x)) - \int (-\cos(x)) e^x dx \\ &= -e^x \cos(x) + \int e^x \cos(x) dx \end{aligned}$$

Hmmm . . . it looks like we'll have to use Integration by Parts again. The "Rule of Thumb" when performing Integration by Parts multiple times is that we don't switch the roles of u and dv . (e.g., if u is an exponential the first time, then u should be the exponential the second time. If dv is the trig function the first time, then dv should be the trig function the second time. Consequently, we have:

u	$= e^x$	dv	$= \cos(x) dx$
$\frac{du}{dx}$	$= e^x$	$\int dv$	$= \int \cos(x) dx$
du	$= e^x dx$	v	$= \sin(x)$

Thus:

$$\begin{aligned} \int e^x \sin(x) dx &= -e^x \cos(x) + \int \underbrace{e^x}_u \underbrace{\cos(x) dx}_{dv} = -e^x \cos(x) + \int u dv \\ &= -e^x \cos(x) + uv - \int v du = -e^x \cos(x) + e^x \sin(x) - \int \sin(x) e^x dx \\ &= -e^x \cos(x) + e^x \sin(x) - \int e^x \sin(x) dx \end{aligned}$$

We have established that:

$$\int e^x \sin(x) dx = -e^x \cos(x) + e^x \sin(x) - \int e^x \sin(x) dx$$

We solve for $\int e^x \cos(x) dx$ algebraically:

$$2 \int e^x \sin(x) dx = -e^x \cos(x) + e^x \sin(x)$$

$$\Rightarrow \int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x) + C$$

i.e., $\int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x) + C$
--

WOW! Extra (10 pts - all or nothing)

$$\int x \arcsin(x) dx =$$

$\arcsin(x)$ is a transcendental function whose derivative is algebraic. $\left(\frac{d}{dx} [\arcsin(x)] = \frac{1}{\sqrt{1-x^2}}\right)$

In such a case, we let the transcendental function be u .

u	$= \arcsin(x)$	dv	$= x dx$
$\frac{du}{dx}$	$= \frac{1}{\sqrt{1-x^2}}$	$\int dv$	$= \int x dx$
du	$= \frac{1}{\sqrt{1-x^2}} dx$	v	$= \frac{1}{2} x^2$

$$\begin{aligned} \text{Therefore, } \int \underbrace{\arcsin(x)}_u \underbrace{x dx}_{dv} &= \int u dv = uv - \int v du = \arcsin(x) \left(\frac{1}{2}x^2\right) - \int \left(\frac{1}{2}x^2\right) \frac{1}{\sqrt{1-x^2}} dx \\ &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{x^2}{\sqrt{1-x^2}} dx \end{aligned}$$

We have: $\int x \arcsin(x) dx = \dots = \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{x^2}{\sqrt{1-x^2}} dx$


x^2	$= \sin^2(\theta)$
x	$= \sin(\theta)$
$\frac{dx}{d\theta}$	$= \cos(\theta)$
dx	$= \cos(\theta) d\theta$

This substitution yields:

$$\begin{aligned} \int x \arcsin(x) dx &= \dots = \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{\sin^2(\theta)}{\sqrt{1-\sin^2(\theta)}} \cos(\theta) d\theta \\ &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{\sin^2(\theta)}{\sqrt{\cos^2(\theta)}} \cos(\theta) d\theta = \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{\sin^2(\theta)}{\cos(\theta)} \cos(\theta) d\theta \\ &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \sin^2(\theta) d\theta = \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{1-\cos(2\theta)}{2} d\theta \\ &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \int (1 - \cos(2\theta)) d\theta \\ &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \left(\theta - \frac{1}{2} \sin(2\theta)\right) + C \end{aligned}$$

Recall: $x = \sin(\theta) \Rightarrow \theta = \arcsin(x)$

The triangle below depicts this relationship:



We have:

$$\begin{aligned}
 \int x \arcsin(x) dx &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \left(\theta - \frac{1}{2} \sin(2\theta) \right) + C \\
 &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \arcsin(x) - \frac{1}{8} (2 \sin(\theta) \cos(\theta)) + C \\
 &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \arcsin(x) - \frac{1}{4} \sin(\theta) \cos(\theta) + C \\
 &= \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \arcsin(x) - \frac{1}{4}x\sqrt{1-x^2} + C
 \end{aligned}$$

$$\int x \arcsin(x) dx = \frac{1}{2}x^2 \arcsin(x) - \frac{1}{4} \arcsin(x) - \frac{1}{4}x\sqrt{1-x^2} + C$$

Note: From the triangle, we used the facts that:

$$\sin(\theta) = \frac{\text{opp}}{\text{hyp}} = \frac{x}{1} = x$$

and

$$\cos(\theta) = \frac{\text{adj}}{\text{hyp}} = \frac{\sqrt{1-x^2}}{1} = \sqrt{1-x^2}$$