



3. Find the area bounded by the graphs of  $f(x) = 4 - x^2$  and  $g(x) = -2x + 4$ . (Partition the appropriate interval, sketch the  $i^{\text{th}}$  rectangle, build the Riemann Sum, derive the appropriate integral.)

4. Five pounds of force is required to stretch a spring 3 inches past the point of equilibrium. How much work is done stretching the spring 12 inches past the point of equilibrium? (Partition the appropriate interval, compute  $F_i$ , build the Riemann Sum, derive the appropriate integral.)

5. Use the “disc method” to compute the volume of the solid of revolution generated by revolving the region bounded by the graphs of  $f(x) = \frac{1}{2}x$ ,  $x = 1$ ,  $x = 3$ , and the  $x$ -axis, about the  $x$ -axis. (For your information: the equation of the  $x$ -axis is  $y = 0$ .)

Use the “five step method” (partition the interval, sketch the  $i^{\text{th}}$  rectangle, form the sum, take the limit)

6. Use the “shell method” to compute the volume of the solid of revolution generated by revolving the region described below about the line  $x = -1$ .

The region lies to the right of the  $y$ -axis and is bounded by the graph  $f(x) = x^2 + 1$ , the  $y$ -axis, and the line  $y = 5$ .

Use the “five step method” (partition the interval, sketch the  $i^{\text{th}}$  rectangle, form the sum, take the limit)